

Quantum Quenches in Extended Systems

John Cardy

University of Oxford

Perugia, July 2007

Joint work with P. Calabrese (Pisa) and S. Sotiriadis (Oxford)

Statement of the problem

- ▶ prepare a system at time $t = 0$ in the ground state $|\psi_0\rangle$ of a (regularised) QFT with hamiltonian H_0
- ▶ for time $t > 0$ evolve *unitarily* with a different hamiltonian H , where $[H, H_0] \neq 0$, e.g. by suddenly changing a parameter – a *quantum quench*, relevant to experiments on cold atoms in optical lattices
- ▶ how do the correlation functions of local operators evolve?
- ▶ for fixed separations, do they become stationary as $t \rightarrow \infty$?
- ▶ do the reduced density matrices of large but finite regions become stationary? If so what is their form?

Simple harmonic oscillator

$$H_0 = \frac{1}{2}p^2 + \frac{1}{2}\omega_0^2 q^2 \qquad H = \frac{1}{2}p^2 + \frac{1}{2}\omega^2 q^2$$

Heisenberg equation of motion has solution

$$q(t) = q(0) \cos \omega t + (p(0)/\omega) \sin \omega t$$

Using $\langle q(0)^2 \rangle = 1/2\omega_0$, $\langle p(0)^2 \rangle = \omega_0/2$,
 $\langle q(0)p(0) + p(0)q(0) \rangle = 0$, and $[q(0), p(0)] = i$ we get the propagator

$$\begin{aligned} \langle T(q(t_1)q(t_2)) \rangle &= \frac{1}{4} \left(\frac{1}{\omega_0} + \frac{\omega_0}{\omega^2} \right) \cos \omega(t_1 - t_2) - \frac{i}{2\omega} \sin \omega|t_1 - t_2| \\ &\quad + \frac{1}{4} \left(\frac{1}{\omega_0} - \frac{\omega_0}{\omega^2} \right) \cos \omega(t_1 + t_2) \end{aligned}$$

Imaginary time


$$\begin{aligned}\langle T(q(\tau_1)q(\tau_2)) \rangle &= \frac{1}{4} \left(\frac{1}{\omega_0} + \frac{\omega_0}{\omega^2} \right) \cosh \omega(\tau_1 - \tau_2) - \frac{2}{\omega} \sinh \omega |\tau_1 - \tau_2| \\ &\quad + \frac{1}{4} \left(\frac{1}{\omega_0} - \frac{\omega_0}{\omega^2} \right) \cosh \omega(\tau_1 + \tau_2)\end{aligned}$$

Imaginary time

$$\begin{aligned}\langle T(q(\tau_1)q(\tau_2)) \rangle &= \frac{1}{4} \left(\frac{1}{\omega_0} + \frac{\omega_0}{\omega^2} \right) \cosh \omega(\tau_1 - \tau_2) - \frac{2}{\omega} \sinh \omega |\tau_1 - \tau_2| \\ &\quad + \frac{1}{4} \left(\frac{1}{\omega_0} - \frac{\omega_0}{\omega^2} \right) \cosh \omega(\tau_1 + \tau_2)\end{aligned}$$


- ▶ $\langle q(\tau)^2 \rangle = 0$ when $\tau = \pm L/2$, where $\omega/\omega_0 = \tanh(L/2)$
- ▶ path integral in imaginary time is the same as if the theory were confined to a slab $-\frac{1}{2}L < \tau < \frac{1}{2}L$ with Dirichlet boundary conditions

Method of Images



- ▶ dependence on $\tau_1 - \tau_2 \leftrightarrow$ (positive) images at $\tau_1 = \tau_2 + 2nL$
- ▶ dependence on $\tau_1 + \tau_2 \leftrightarrow$ (negative) images at $\tau_1 = -\tau_2 + 2nL$

Method of Images



- ▶ dependence on $\tau_1 - \tau_2 \leftrightarrow$ (positive) images at $\tau_1 = \tau_2 + 2nL$
- ▶ dependence on $\tau_1 + \tau_2 \leftrightarrow$ (negative) images at $\tau_1 = -\tau_2 + 2nL$
- ▶ if we ignore (or average over) the oscillating term, the propagator is the *same* as that at finite temperature $\beta_{\text{eff}} = 2L$

Coupled harmonic oscillators (free scalar field theory)

- ▶ a collection of oscillators $H = \int (\frac{1}{2}|\pi_k|^2 + \frac{1}{2}\omega_k^2|\phi_k|^2)d^d k$,
 $\omega_k = (m^2 + k^2)^{1/2}$
- ▶ consider a quench $m_0 \rightarrow m$, with $m_0 > m$
- ▶ the oscillating part in $\langle T(\phi(t_1, x_1)\phi(t_2, x_2)) \rangle$ has the form

$$\int \frac{d^d k}{(2\pi)^d} e^{ik(x_1 - x_2)} \left(\frac{1}{\omega_{0k}} - \frac{\omega_{0k}}{\omega_k^2} \right) \cos(\omega_k(t_1 + t_2))$$

- ▶ if $\omega_k = (k^2 + m^2)^{1/2}$ with $m > 0$ the second term
 $\sim t_1^{-d/2} \cos(2mt_1) \rightarrow 0$ as $t_1 \sim t_2 \rightarrow \infty$

- ▶ the remainder corresponds to an effective k -dependent temperature

$$\beta_k = (4/\omega_k) \tanh^{-1} (\omega_k/\omega_{0k})$$

- ▶ if $|x_1 - x_2| \ll t$ the dominant contribution comes from $k \sim 0$, and we can ignore the k -dependence in β_k
- ▶ the 2-point function (and the N -point functions) all thermalize (but slowly)

Onset of correlations

- for large m_0

$$\langle \phi(t_1, x_1) \phi(t_2, x_2) \rangle \sim m_0 \int d^d k \frac{e^{ik(x_1 - x_2)}}{\omega_k^2} (\cos \omega_k(t_1 - t_2) - \cos \omega_k(t_1 + t_2))$$

$$\begin{aligned} \frac{\partial}{\partial t_1}(\text{this}) \propto & G_F(x_1 - x_2, t_1 - t_2) - G_F(x_1 - x_2, -t_1 + t_2) \\ & - G_F(x_1 - x_2, t_1 + t_2) + G_F(x_1 - x_2, -t_1 - t_2) \end{aligned}$$

- if $t_1 + t_2 < |x_1 - x_2|$ this vanishes by Lorentz invariance – *horizon effect*
- in general behaviour near horizon is smoothed out over scales $\delta t \sim m_0^{-1}$

Massless case (conformal field theory)

- ▶ for $m = 0$ in 1+1 dimensions we find instead

$$\langle \phi(t_1, x_1) \phi(t_2, x_2) \rangle = \begin{cases} 0 & \text{if } t_1 + t_2 < |x_1 - x_2| \\ m_0(t_1 + t_2 - |x_1 - x_2|) & \text{if } t_1 + t_2 > |x_1 - x_2| \end{cases}$$

- ▶ many gapless interacting systems in $d = 1$ are equivalent to *conformal field theories*
- ▶ local observables

$$\Phi_q(x, t) \sim e^{iq\phi(x, t)}$$

where $\phi(x, t)$ is a massless free field

Massless case (conformal field theory)

- ▶ for $m = 0$ in 1+1 dimensions we find instead

$$\langle \phi(t_1, x_1) \phi(t_2, x_2) \rangle = \begin{cases} 0 & \text{if } t_1 + t_2 < |x_1 - x_2| \\ m_0(t_1 + t_2 - |x_1 - x_2|) & \text{if } t_1 + t_2 > |x_1 - x_2| \end{cases}$$

- ▶ many gapless interacting systems in $d = 1$ are equivalent to *conformal field theories*
- ▶ local observables

$$\Phi_q(x, t) \sim e^{iq\phi(x, t)}$$

where $\phi(x, t)$ is a massless free field

1-point functions:

$$\langle \Phi_q(x, t) \rangle = e^{-(q^2/2)\langle \phi(x, t)^2 \rangle} \sim e^{-m_0 q^2 t}$$

2-point functions:

$$\langle \Phi_q(x_1, t_1) \Phi_{-q}(x_2, t_2) \rangle = e^{-(q^2/2) \langle (\phi(x_1, t_1) - \phi(x_2, t_2))^2 \rangle}$$

so, for $t_1 + t_2 < |x_1 - x_2|/c$,

$$\langle \Phi_q(x_1, t_1) \Phi_{-q}(x_2, t_2) \rangle \sim \langle \Phi_q(x_1, t_1) \rangle \langle \Phi_{-q}(x_2, t_2) \rangle$$


while for $t_1 + t_2 > |x_1 - x_2|/c$

$$\langle \Phi_q(x_1, t_1) \Phi_{-q}(x_2, t_2) \rangle \sim e^{-m_0 q^2 (t_1 + t_2 - (t_1 + t_2 - |x_1 - x_2|/c))} = e^{-m_0 q^2 |x_1 - x_2|/c}$$


- thermalization of a region of length ℓ takes place exponentially fast (on a time-scale $O(m_0^{-1})$) after the end-points come into mutual causal contact
- these results hold for any CFT in 1+1 dimensions

Physical picture

- ▶ $|\psi_0\rangle$ has (extensively) higher energy than the ground state of H
- ▶ it acts as a source of (quasi)particles at $t = 0$
- ▶ particles emitted from regions size $\sim m_0^{-1}$ are entangled
- ▶ subsequently they move classically (at velocity $\pm c$)
- ▶ incoherent particles arriving at r from well-separated initial points cause relaxation of local observables (except conserved quantities like the energy) to their ground state values:




- *horizon effect*: local observables with separation r become correlated when left- and right-moving particles originating from the same spatial region $\sim m_0^{-1}$ can first reach them:



- if all particles move at unique speed c correlations are then frozen for $t > r/2c$

- *horizon effect*: local observables with separation r become correlated when left- and right-moving particles originating from the same spatial region $\sim m_0^{-1}$ can first reach them:



- if all particles move at unique speed c correlations are then frozen for $t > r/2c$
- entanglement entropy of an interval of length ℓ is extensive, and identical to Gibbs-Boltzmann entropy at temperature β_{eff}^{-1}



General dispersion relation

$$(\partial/\partial t)\langle\phi(x_1,t)\phi(x_2,t)\rangle = m_0 \int \frac{d^d k e^{ik(x_1-x_2)}}{\omega_k} \sin(2\omega_k t)$$

- ▶ large x, t behaviour given by stationary phase approximation

$$|x_1 - x_2|/2t = d\omega_k/dk = \text{group velocity } v_k$$

- ▶ correlations begin to form at $t = |x_1 - x_2|/2v_{\max}$
- ▶ large t behaviour dominated by slowest moving particles:
eg lattice dispersion relation gives a power law approach to asymptotic limit



- ▶ agrees with exact results for Ising and XY spin chains

General interacting QFTs

- ▶ can we safely ignore the oscillating terms in the propagator within loops?
- ▶ *eg* $\lambda\phi^4$ theory in the Hartree (large N) approximation

The diagram shows a Feynman diagram equation. On the left is a single horizontal line representing a propagator. This is followed by an equals sign. To the right of the equals sign are two terms added together. The first term is another single horizontal line. The second term is a horizontal line with a circle (representing a loop) attached to its midpoint, forming a self-energy loop.

- ▶ even if the renormalized mass is zero, the interactions + the modified propagator generate an effective mass, which means that oscillating terms in the loop are damped \rightarrow thermalisation

Summary and further remarks

- ▶ quantum quenches from $m_0 \downarrow m$ appear to lead to thermalisation of finite regions if $m > 0$, and even when $m = 0$ in the presence of interactions
- ▶ there is a ‘horizon’ effect: correlations only begin to change after points come into mutual causal contact
- ▶ many interesting questions remain: *eg* quenches from a disordered phase \rightarrow ordered phase – can we drive a phase transition by changing the initial state? ...